

Fig 2. Experimental and simulated absorption spectra of L (left); 2D potential energy curves for the ground and first singlet excited states of L<sub>2</sub> with variation of N<sub>1</sub>-H<sub>1</sub> and N<sub>2</sub>-H<sub>2</sub> bond lengths (right)

### References

1. Gladysiak A., Nguyen T. N., Bounds R. et al. // Chem. Sci. 2019. Vol. 10. P. 6140–6148.
2. Bai G., Tsang M.-K., Hao J. // Adv. Funct. Mater. 2016. Vol. 26. P. 6330–6350.
3. Shi C., Shen X., Zhu Y. et al. // ACS Appl. Mater. Interfaces, 2019. Vol. 11. P. 18548–18554.
4. Waluk J. // Journal of Molecular Structure. 1988. Vol. 177. P. 415–419.
5. Serdiuk I. E., Roshal A. D. // Dyes and Pigments. 2017. Vol. 138. P. 223–244.
6. Vinogradova K. A., Shekhovtsov N. A., Berezin A. S. et al. // Inorg. Chem. Comm. 2019. Vol. 100. P. 11–15.

\* The research was supported by Russian Foundation for Basic Research and the government of the Novosibirsk region of the Russian Federation, grant № 18-43-540016 r\_a.

УДК 547.745

E. E. Stepanova, A. N. Maslivets

Perm State University,  
614990, Russia, Perm, Bukireva St., 15,  
caterina.stepanova@psu.ru

## 1,3-DIPOLAR CYCLOADDITION REACTION OF FUSED 1H-PYRROLE-2,3-DIONES WITH NITRONES: A DIVERGENT APPROACH TO PYRROLOISOXAZOLES\*


**Keywords:** 1,3-dipolar cycloaddition, isoxazole, nitrones, 1H-pyrrole-2,3-dione, rearrangement.

Divergent synthesis is a promising trend in small molecules drug discovery [1, 2]. It allows to investigate the chemical space more efficiently and economically since diverse compounds collections are created from a limited set of reagents.

1,3-Dipolar cycloaddition is a facile tool for simultaneous introduction of several stereogenic centers through a single step to afford complex heterocyclic structures [3]. In the view of requirements of green chemistry, it seems to be a beneficial synthetic approach since it proceeds in an atom and step economic manner along with high regio- and stereoselectivity.

1*H*-Pyrrole-2,3-diones fused at [*e*]-side (hetareno[*e*]pyrrole-2,3-diones) are readily available polyelectrophilic reagents enabling synthesis of various heterocyclic systems with divergent skeletons [4, 5].

We have developed a divergent approach to skeletally diverse isoxazoles *via* 1,3-dipolar cycloaddition reaction of hetareno[*e*]pyrrole-2,3-diones **1** with nitrones **2**. Isoxazoles **3** [6] and **4** can be obtained from a single set of reagents (compounds **1** and **2**), and the formation of compounds **3** or **4** proceeds regioselectively in dependence on the reaction conditions (solvent and temperature).



## References

1. Galloway W. R., Isidro-Llobet A., Spring D. R. // *Nature Communications*. 2010. Vol. 1, № 80.
2. Garcia-Castro M., Zimmermann S., Sankar M. G. et al. // *Angewandte Chemie International Edition*. 2016. Vol. 55. P. 7586–7605.
3. Hashimoto T., Maruoka K. // *Chemical Reviews*. 2015. Vol. 115. P. 5366–5412.
4. Konovalova V. V., Shklyayev Yu. V., Maslivets A. N. // *Arkivoc*. 2015. Vol. I. P. 48–69.
5. Konovalova V., Maslivets A. // *Mini-Reviews in Organic Chemistry*. 2018. Vol. 16. P. 173–192.
6. Stepanova E. E., Dmitriev M. V., Maslivets A. N. // *Tetrahedron Letters*. 2020. Vol. 56. № 151595.

\* This work was supported by the Russian Science Foundation, project № 19-13-00290.